

Comment on “No enhancement of the localization length for two interacting particles in a random potential”

In a recent letter [1] Römer and Schreiber report on numerical calculations that led them to conclude that the previously observed enhancement [2–6] of the localization length L_2 of two interacting particles (TIP) vanishes in the thermodynamic limit. Such a claim (i) is in conflict with the scaling theory of localization, (ii) ignores a consistent picture from a wealth of published numerical data and analytical investigations, and (iii) directly contradicts new numerical results obtained with a Green function method [6] that is well adapted to the problem under study.

(i) Römer and Schreiber considered system sizes L between 80 and 360 for their extrapolation. For $L = 360$ they find $L_2 \approx 18$, i.e. L_2 exceeds the one-particle localization length $L_1 \approx 12$ significantly. How can these extended states shrink upon increasing the system size from $20L_2$ to, say, $25L_2$? According to Thouless’ scaling picture, the energy shift associated with such an increase of L amounts to a negligible fraction of the mean level spacing, and no significant effect can be expected.

(ii) Let n_1 and n_2 be the sites of the two particles. Shepelyansky showed [2] that, for a small fraction ($\approx L_1/(2L)$) of states, the center of mass $(n_1 + n_2)/2$ is extended over L_2 , while $|n_1 - n_2| \lesssim L_1$. Alternative and complementary derivations are provided by Imry’s variant [3] of the Thouless scaling picture, and by an effective σ model [7], both not restricted to finite L . Three different numerical methods have confirmed this effect. First, extended states of the type described above were obtained by exact diagonalization [5] of the TIP Hamiltonian in a ring geometry. Second, the delocalization of the center of mass was studied in Ref. [6] by calculating the TIP Green function $\langle n_1 n_2 | G_2(E) | n'_1 n'_2 \rangle$ for $n_1 = n_2 = 0$ and $n'_1 = n'_2 = L$, i.e. *pair transfer* from 0 to L . Third, the transport of *one* particle through a relatively small sample containing the second particle was investigated [4] with the transfer matrix method. They all reach a common conclusion: For an on-site interaction $U = 1$ and $L_1 \approx 11$, one has $L_2 \approx 25 > L_1$. Comparable results were obtained in *infinite* systems [4] with a prescribed maximum separation $|n_1 - n_2|$ (“bag model”).

(iii) Römer and Schreiber used transfer matrix methods (similar to those in Ref. [4]), and concluded that $L_2 \rightarrow L_1$ when $L \rightarrow \infty$. In Fig. 1, we compare data from Ref. [1] with results from the Green function method ($U = 1, L_1 \approx 11$). The latter consistently gives $L_2 \approx 25 \approx 2L_1$ up to the maximum size we considered, $L = 1000$. Without explaining the data of Ref. [1] in detail, we emphasize the following important problem: The L^2 entries in the transfer matrix correspond to the L^2 functions $\langle 0 n_2 | G_2(E) | L n'_2 \rangle$. Most of these characterize single particle transport (decay scale L_1), only those

with $n_2 \lesssim L_1, n'_2 \gtrsim L - L_1$ describe pair transport (decay scale $L_2/2$ [6], with the conventions of Ref. [1]). To distinguish these two scales one has to either suppress single particle transport (Green function method, bag model) or ensure that $L_2 \gg 2L_1$ (as in Ref. [4]). Unfortunately, Ref. [1] meets neither of these requirements, and it is unclear, which scale is actually measured.

In conclusion, the central claim in Ref. [1] is due to a serious misinterpretation, and the localization length L_2 of the interaction-assisted states is *independent* of L , as expected on physical grounds.




FIG. 1. Data for L_2 from Ref. [1] (squares) and the Green function method (circles) versus $L^{-1/2}$. We thank F. v. Oppen and T. Wettig for their computer program.

Klaus Frahm¹, Axel Müller-Groeling², Jean-Louis Pichard³, and Dietmar Weinmann⁴

¹Laboratoire de Physique Quantique, Université Paul Sabatier, F-31062 Toulouse, France

²Max-Planck-Institut für Kernphysik, D-69029 Heidelberg, Germany

³SPEC, CEA-Saclay, F-91191 Gif-sur-Yvette, France

⁴Institut für Physik, Universität Augsburg, D-86135 Augsburg, Germany

PACS numbers: 72.15.Rn, 71.55.Jv, 72.10.Bg

- [1] R.A. Römer and M. Schreiber, Phys. Rev. Lett. **78**, 515 (1997).
- [2] D. L. Shepelyansky, Phys. Rev. Lett. **73**, 2607 (1994).
- [3] Y. Imry, Europhys. Lett. **30**, 405 (1995).
- [4] K. Frahm, A. Müller-Groeling, J.-L. Pichard, and D. Weinmann, Europhys. Lett. **31**, 169 (1995).
- [5] D. Weinmann, A. Müller-Groeling, J.-L. Pichard, and K. Frahm, Phys. Rev. Lett. **75**, 1598 (1995).
- [6] F. von Oppen, T. Wettig, and J. Müller, Phys. Rev. Lett. **76**, 491 (1996).
- [7] K. Frahm, A. Müller-Groeling, and J.-L. Pichard, Phys. Rev. Lett. **76**, 1509 (1996).